April 2, 1997
hep-ph/9704219

UICHEP-TH/97-2

arXiv:hep-ph/9704219v2 8 Apr 1997

CP Violation and Scalar Lepton Flavor Oscillation

David Bowser-Chao and Wai-Yee Keung

Physics Department, University of Illinois at Chicago, IL 60607-7059, USA

Abstract

Lepton flavor violation can be induced in supersymmetry by the mixing of two or more of the leptonic scalar partners, $\tilde{l} \to \tilde{l}'$, where $\tilde{l}, \tilde{l}' = \tilde{e}, \tilde{\mu}, \tilde{\tau}$. Krashnikov and Arkani-Hamed *et al.* pointed out that this effect may be observable at the Next Linear Collider, through slepton pair production with subsequent lepton flavor violating decays. If the slepton mixing involves all three generations, CP violation from the Cabibbo-Kobayashi-Maskawa phase could lead to asymmetries between the observed LFV decays $\tilde{l} \to \tilde{l}'$ and $\bar{\tilde{l}} \to \bar{\tilde{l}}'$. We lay down the formalism and give simple expressions for the CP violating asymmetry in the transition probabilities, and consider possible signals at future colliders.

PACS numbers: 11.30.Er, 11.30.Hv, 14.80.Ly, 14.60.Gh

Introduction

Quark flavor is not conserved in nature, as manifestly demonstrated by oscillation in the K–$\bar{K}$ system. There is no evidence to date, however, for the analogous lepton flavor violation. The branching ratio for $\mu \to e\gamma$, for example, is less than 4.9×10^{-11} at the 95% confidence level[1]. This is well in accord with the Standard Model, where lepton flavor violation (LFV) is expected to be highly suppressed, due to the smallness of m_ν/v, where m_ν is the largest neutrino mass and v the electroweak breaking scale.

Lepton flavor violation is also suppressed in exactly supersymmetric extensions of the Standard Model; supersymmetry, however, is necessarily broken, and soft symmetry breaking terms can in general generate significant LFV[2]. Recently, Krashnikov[3] and Arkani-Hamed[4] *et al.* showed that the resultant mixing between at least two of the charged slepton generations could produce dramatic lepton flavor violating signals at LEPII or the Next Linear Collider (NLC). While detailing the particular case of two generation mixing, these authors also noted the obvious extension to mixing amongst all three generations.

Beyond simply complicating the analysis of lepton flavor violation, moving to three generation mixing also gives rise to a new source of CP violation (CPV); namely, the single complex phase in the Cabibbo-Kobayashi-Maskawa (CKM) matrix that relates the electroweak flavor basis to the three mass eigenstates. In this paper we examine the consequences of this CP violating phase[5].

The analysis is similar in spirit to CPV from three family neutrino oscillation[6]. We shall provide compact formulas for the CPV transition probability asymmetry $P(\tilde{\bar{l}} \to \bar{l}') - P(\tilde{l} \to l')$, where $\tilde{l}, \tilde{l}'$ denote the sleptons $\tilde{e}, \tilde{\mu}, \tilde{\tau}$ and $\tilde{\bar{l}}, \tilde{\bar{l}}'$ are the respective anti-particles. Following Ref.[6], we show that the CPV result is identical for the available choices $(l, l') = (e, \mu), (\mu, \tau)$, and (τ, e) , then consider observation of the asymmetry at future colliders.

Formalism

First, we shall assume (as did Refs. [3, 4]) (1) the presence of LFV soft symmetry breaking terms in the electroweak scale lagrangian, without regard for the nature of their origin, and (2) that left-right slepton mixing is small. The second assumption simplifies analysis of CPV and LPV effects, which then have separate contributions from the left and right-handed sleptons. This assumption is not strongly model-dependent, since left-right slepton mixing is generally suppressed by the ratio of the lepton mass to the left-right slepton mass splitting, with the latter set by the D-term to be of order the electroweak scale[7]. Furthermore, we shall see below that CPV effects involving left-right mixing are further suppressed by the the ratio of the slepton width to the left-right mass splitting. We shall thus follow Refs. [3, 4] and focus on the right-handed sleptons, while noting that generalization of our analysis to the separate left-handed slepton contribution to CPV is also possible.

Due to mixing, the flavor states ($\tilde{e}$, $\tilde{\mu}$, $\tilde{\tau}$) do not evolve in time with a trivial phase. We define the mixing amplitudes $U_{\alpha i}$, where the flavor state is expressed by superposition of mass eigenstates $|S_i\rangle$ of masses m_i,

$$|\tilde{l}_\alpha\rangle = \sum_i U_{\alpha i}|S_i\rangle \ , \quad |S_i\rangle = \sum_i U^*_{\alpha i}|\tilde{l}_\alpha\rangle \ . \tag{1}$$

The matrix U can be explicitly written using the standard KM parameterization with three mixing angles θ_i and one phase δ,

$$\begin{pmatrix} \tilde{e} \\ \tilde{\mu} \\ \tilde{\tau} \end{pmatrix} = \begin{pmatrix} c_1 & s_1c_3 & s_1s_3 \\ -s_1c_2 & c_1c_2c_3 + s_2s_3e^{i\delta} & c_1c_2s_3 - s_2c_3e^{i\delta} \\ -s_1s_2 & c_1s_2c_3 - c_2s_3e^{i\delta} & c_1s_2s_3 + c_2c_3e^{i\delta} \end{pmatrix} \begin{pmatrix} S_1 \\ S_2 \\ S_3 \end{pmatrix} \ , \tag{2}$$

with $s_i = \sin\theta_i$ and $c_i = \cos\theta_i$.

If the LSP is a neutralino χ^0 (which we assume to be primarily Bino), the flavor eigenstate l can be tagged by the decay $\tilde{l} \to l\chi^0$. The flavor state $|\tilde{\alpha}\rangle$ ($\alpha = e, \mu$, or τ) of a slepton produced at $t = 0$ in the rest frame, could be resolved as $|\tilde{\beta}\rangle$ at a later time t with the transition amplitude,

$$A(\tilde{\alpha} \to \tilde{\beta}; t) = \sum_i U_{\alpha i} U^*_{\beta i} \exp(-im_i t - \tfrac{1}{2}\Gamma t) \ . \tag{3}$$

For simplicity we assume this to be the only open decay channel; if not, results below should be multiplied by the decay branching ratios[4]. We also take all states to have a common decay width Γ and

$$\Delta m_{ij} \equiv m_i - m_j \ ; \ \Delta m_{ij}, \Gamma \ll \bar{m} \equiv \tfrac{1}{3}(m_1 + m_2 + m_3) \ . \tag{4}$$

Further restrictions on Δm_{ij} are discussed below.

The time-dependent oscillation probability is $p(\tilde{\alpha} \to \beta; t) = |A(\tilde{\alpha} \to \tilde{\beta}; t)|^2$; note that we omit the decay product neutralino in our notation. The time–averaged probability that is actually measured is given by

$$\begin{aligned} P(\tilde{\alpha} \to \beta) &= \int_0^\infty p(\tilde{\alpha} \to \tilde{\beta}; t) dt \Big/ \int_0^\infty \sum_\beta p(\tilde{\alpha} \to \tilde{\beta}; t) dt \\ &= U_{\alpha i} U^*_{\beta i} U^*_{\alpha j} U_{\beta j} \left(\frac{\Gamma}{\Gamma + (m_i - m_j) i} \right) = P(\tilde{\bar{\beta}} \to \bar{\alpha}) \ . \end{aligned} \tag{5}$$

From the unitarity of $U_{\alpha i}$ and the second line of the equation above, it can be seen that lepton flavor violation vanishes[4] as $\Delta m_{ij} \ll \Gamma$; in the limit of exactly degenerate sleptons, the mass basis can be arbitrarily rotated to bring $U_{\alpha i}$ into diagonal form.

CP violation is reflected in asymmetry between the transition probabilities of CP–conjugate channels, $P(\alpha \to \beta) - P(\tilde{\bar{\alpha}} \to \bar{\beta})$. Since $P(\tilde{\bar{\alpha}} \to \bar{\beta}) = P(\tilde{\beta} \to \alpha)$ from CPT invariance, the difference flips sign under interchange of α and β and vanishes for the diagonal case $\alpha = \beta$. It is convenient to introduce the antisymmetric symbol $\varepsilon_{\alpha\beta}$ such that $\varepsilon_{e\mu} = \varepsilon_{\mu\tau} = \varepsilon_{\tau e} = 1$. We summarize all properties of asymmetry,

$$P(\tilde{\alpha} \to \beta) - P(\tilde{\bar{\alpha}} \to \bar{\beta}) = \varepsilon_{\alpha\beta} \mathcal{A} \ , \quad \mathcal{A} = -4 \ \mathrm{Im}(U_{e1} U^*_{\mu 1} U^*_{e2} U_{\mu 2}) \ \mathrm{Im}(\phi_{12} + \phi_{23} + \phi_{31}) \ , \tag{6}$$

where

$$\phi_{ij} = \frac{\Gamma}{\Gamma + (m_i - m_j) i} \ , \quad \mathrm{Im}\phi_{ij} = \frac{(m_j - m_i)\Gamma}{\Gamma^2 + (m_i - m_j)^2} \ . \tag{7}$$

From Eq.(4), this asymmetry result could also have been obtained by using the narrow width approximation. Note that $\mathcal{A}$ vanishes not just for $\Delta m_{ij} \ll \Gamma$ as discussed above, but also if $\Delta m_{ij} \gg \Gamma$, since in this limit $\mathrm{Im}\phi_{ij}$ vanishes as well. This is simply due to the fact that CP violation requires interference between $U_{\alpha i}$ and the final state decay phases in in Eq.(5); in the limit of very large slepton mass splitting, Eq.(5) becomes an incoherent sum over the mass eigenstates. For CP violation from slepton mixing, we thus require $\Delta m_{ij} \sim \Gamma$. We can also see that CPV effects involving left-right mixing are indeed suppressed by the the ratio of the slepton width to the left-right mass splitting.

The expression for $\mathcal{A}$ involves the single universal CPV combination[8] of mixing amplitudes in the case of three families,

$$X_{\rm CP} \equiv {\rm Im}(U_{e1}U^*_{\mu 1}U^*_{e2}U_{\mu 2}) = -s_1^2 c_1 s_2 c_2 s_3 c_3 \sin\delta \ . \tag{8}$$

The magnitude of this CPV parameter $X_{\rm CP}$ ranges from zero to the maximum value $\sqrt{3}/18 = 0.096$. Ref.[4] shows that the present experimental bound on ${\rm Br}(\mu \to e\gamma)$ puts no significant constraint on the mixing angles, with $(\Delta m/\bar{m}) \sin 2\theta \lesssim 0.01$.

One may wonder how the experimental limit[9] on the electric dipole moment (EDM) of the electron constrains the phase in U. It turns out that at the one loop level, the EDM requires both the left and right handed sleptons to participate in the loop. Each handedness contributes a GIM-like suppression factor, $\Delta m_{L,R}/\bar{m}_{L,R}$($\approx$ 0.01 for our case). Together with a factor $m_l/\bar{m}$ from the soft SUSY-breaking A_l term or the supersymmetric μ term, which mixes the L and R components, there is sufficient suppression to expect no significant constraint on the phase (note that the electron EDM *does* constrain the phase in the left-right slepton mixing[2], as well as putting tight limits on a relative phase between the complex mass of the gauge fermion and the A_l or μ term[10]).

Experiments

It is believed that the cleanest environment to search for sleptons is at the proposed Next Linear e^+e^- collider (NLC) at an energy of $\sqrt{s} \simeq 500$ GeV to 1.5 TeV. The signature for pair production of sleptons is, assuming the neutralino decay discussed above, a pair of charged leptons plus missing energy. In Fig. 1, we illustrate the cross section for right handed slepton production. At $\sqrt{s} = 500$ GeV, $\sigma(\tilde{e}\tilde{\bar{e}})$ is almost as large as 1 pb under the assumption that the neutralino exchanged in the t-channel is purely Bino with mass $M_\chi = 50$ GeV (with an obvious degradation for an admixture of Bino). Ref. [4] calculated the backgrounds from WW, $e^\pm \nu W^\mp$, and $(e^+e^-)W^+W^-$ to total about 12 pb. Through efficient cuts[11], Ref. [4] estimates a reduction of the background to about 5 fb for unpolarized beams, with 30%

signal acceptance, with further improvement possible with right-handed polarized beams. Because the CPV signals considered here are simply asymmetries in the LFV signals of the sort considered in Ref. [4], we shall simply rely on these figures to provide an estimate of the sensitivity to CP violation from slepton mixing.

We start with a simple CP-odd observable which ignores the full correlation between the lepton l and the anti-lepton $\bar{l}'$. From the sample of dilepton events that pass the cuts, we consider the CPV asymmetry between muon and anti-muon events,

$$S_\mu = \frac{N(\mu^-) - N(\mu^+)}{N(\text{all signal})} \ . \tag{9}$$

Clearly only unequal flavor dilepton ($\mu^\pm l^\mp, l \neq \mu$) events contribute to this asymmetry.

Since S_μ only involves a single particle count, it can be expressed directly in terms of the single particle transition probability (Eq.(5)),

$$S_\mu = \frac{\sigma(\tilde{e}\tilde{\bar{e}})[P(\tilde{e} \to \mu) - P(\tilde{\bar{e}} \to \bar{\mu})] + \sigma(\tilde{\tau}\tilde{\bar{\tau}})[P(\tilde{\tau} \to \mu) - P(\tilde{\bar{\tau}} \to \bar{\mu})]}{\sigma(\tilde{e}\tilde{\bar{e}}) + \sigma(\tilde{\mu}\tilde{\bar{\mu}}) + \sigma(\tilde{\tau}\tilde{\bar{\tau}})} \ . \tag{10}$$

Using Eq. (6), we obtain

$$S_\mu = \mathcal{A} f \ , \quad f \equiv \frac{\sigma(\tilde{e}\tilde{\bar{e}}) - \sigma(\tilde{\tau}\tilde{\bar{\tau}})}{\sigma(\tilde{e}\tilde{\bar{e}}) + \sigma(\tilde{\mu}\tilde{\bar{\mu}}) + \sigma(\tilde{\tau}\tilde{\bar{\tau}})} \ . \tag{11}$$

At the tree level, $\sigma(\tilde{\tau}\tilde{\bar{\tau}})$ and $\sigma(\tilde{\mu}\tilde{\bar{\mu}})$ come only from the γ-Z exchange amplitude in the s channel, but $\sigma(\tilde{e}\tilde{\bar{e}})$ involves additional neutralino exchange amplitude in the t-channel. Thus f in Eq. (11) is generally non-zero. The fraction f can be close to one in the limit that the t-channel diagram dominates, which will of course be true given sufficiently high collider energy. Fig. 1 shows f to be 0.75 at $\sqrt{s} = 500$ GeV and $\bar{m}$=150 GeV, under the assumption that the exchange gaugino is purely Bino with mass $M_\chi = 50$ GeV. With an integrated luminosity of 100 fb^{-1}, around at least 10^4 events from the slepton pair production will survive the dedicated cuts[4, 11] which severely reduce the predictable backgrounds to only about 500 $\mu^\mp e^\pm$ events and about the same number of $\mu^\mp \tau^\pm$ events. Thus, it is possible to measure the CP asymmetry S_μ at the level 0.02 with 5σ.

The other flavor asymmetries are $S_e = 0$ (which vanishes because $\sigma(\tilde{\tau}\tilde{\bar{\tau}}) = \sigma(\tilde{\mu}\tilde{\bar{\mu}})$) and $S_\mu = -S_\tau$. Statistics will be doubled if we take the combination $S_\mu - S_\tau$.

In Fig. 2, we show the size of $S_\mu - S_\tau$ versus $\Delta m/\Gamma$ for $f = 0.75$, in the scenario that the mass differences are equal in magnitude to the width, namely, $m_1 - m_2 = m_2 - m_3 = \Delta m \sim \Gamma$.

In the scenario where $\sigma(\tilde{e}\bar{\tilde{e}}) \approx \sigma(\tilde{\mu}\bar{\tilde{\mu}}) \approx \sigma(\tilde{\tau}\bar{\tilde{\tau}})$ (e.g., if the t-channel gaugino mass is relatively high), we can no longer use the uncorrelated asymmetry S to probe CPV. We can, however, turn to other correlated observables such as $S_{\mu\bar{e}}$, which is defined by replacing the numerator in Eq.(9) by $N(\mu^- e^+) - N(\mu^+ e^-)$.

The overall amplitude for tagging $\mu\bar{e}$ at time t and $\bar{t}$ after the production comprises two contributions. One is the t-channel gaugino exchange amplitude (for the *selectron* production only) which has the time evolving factor

$$A(\tilde{e} \to \mu)A(\bar{\tilde{e}} \to \bar{e}) = \sum_i U_{ei}U^*_{\mu i} \exp(-im_i t - \tfrac{1}{2}\Gamma t) \sum_j U^*_{ej}U_{ej} \exp(-im_j \bar{t} - \tfrac{1}{2}\Gamma \bar{t}) \ . \tag{12}$$

The other is the common s-channel γ–Z exchange amplitude which has a simpler factor due to unitarity of U matrix,

$$\sum_\alpha A(\tilde{\alpha} \to \mu)A(\bar{\tilde{\alpha}} \to \bar{e}) = \sum_i U^*_{\mu i}U_{ei} \exp[(-im_i - \tfrac{1}{2}\Gamma)(t + \bar{t})] \ . \tag{13}$$

Denoting the asymmetry S^0 in the special case that the s channel dominates, we have the simple expression,

$$S^0_{\mu\bar{e}} = -\tfrac{1}{3} \cdot 4X_{\rm CP}\ {\rm Im}(\phi^2_{12} + \phi^2_{23} + \phi^2_{32}) = -S^0_{\mu\bar{\tau}} = S^0_{e\bar{\tau}} \ . \tag{14}$$

The statistics factor $\frac{1}{3}$ can be offset by summing all three flavor asymmetries, as in $S^0_{\rm tot} \equiv S^0_{\mu\bar{e}} - S^0_{\mu\bar{\tau}} + S^0_{e\bar{\tau}}$.

The asymmetry formula in the general case, where both (s and t) channels are important, is lengthy but quite straightforward, and which we present next.

The amplitudes for $e^-e^+ \to \tilde{l}\,\bar{\tilde{l}}$ consist of a common element M_0 from the s-channel γ–Z exchange and an additional piece M' for the case $l = e$ from the t-channel gaugino exchange. Their explicit expressions at the tree level are provided below. From Eqs. (12,13), we derive

$$\frac{d\sigma(e^-e^+ \to \tilde{l}\,\bar{\tilde{l}} \to \alpha\bar{\beta} + \text{missing neutralinos})}{d\cos\theta} = K\Big(|M_0|^2 \sum_{i,j} U^*_{\alpha i}U_{\beta i}U_{\alpha j}U^*_{\beta j}\phi^2_{ij} \tag{15}$$

$$+|M'|^2 P(\tilde{e} \to \alpha)P(\bar{\tilde{e}} \to \bar{\beta}) + 2{\rm Re}(M_0 M'^* \sum_{ijk} U^*_{\alpha i}U_{\beta i}U^*_{ek}U_{\alpha k}U_{ej}U^*_{\beta j}\phi_{ik}\phi_{ij})\Big) \ .$$

The flux and phase space factor K is given below. The non-zero tree level amplitudes M_0 and M' for different channels are[12]

$$M_0(e^-_L e^+_R \to \tilde{l}_R\,\bar{\tilde{l}}_R) = e^2 s \lambda^{\frac{1}{2}} \sin\theta \left(\frac{1}{s} + \frac{-\frac{1}{2} + x_W}{(1 - x_W)(s - M_Z)}\right) \ , \tag{16}$$

$$M_0(e_R^- e_L^+ \to \tilde{l}_R\ \tilde{\bar{l}}_R) = e^2 s \lambda^{\frac{1}{2}} \sin\theta \left(\frac{1}{s} + \frac{x_W}{(1-x_W)(s-M_Z)} \right) \ , \tag{17}$$

$$M'(e_R^- e_L^+ \to \tilde{e}_R\ \tilde{\bar{e}}_R) = \frac{e^2 s \lambda^{\frac{1}{2}} \sin\theta}{(1-x_W)} \sum_i \frac{|V_{Bi}|^2}{t - M^2_{\chi i}} \ . \tag{18}$$

Here, θ is the CM polar scattering angle, M_{χ_i} are the neutralino masses ($i = 1$ is the lightest), while $|V_{Bi}|^2$ is the mixing probability of the Bino component in the i-th neutralino; $\lambda^{\frac{1}{2}} = \sqrt{1 - 4\bar{m}^2/s}$ is the slepton CM velocity, s and t are the usual invariant squares of energy and momentum transfer, x_W is the electroweak parameter, $M_W^2/M_Z^2 = 1 - x_W$. The flux and phase space factor is $K = \lambda^{\frac{1}{2}}/(128\pi s)$.

The differential rate difference between CP conjugate channels is

$$\frac{d\Delta R_{\alpha\bar{\beta}}}{d\cos\theta} = K\Big(|M_0|^2 \hat{S}^0_{\alpha\bar{\beta}} + |M'|^2 \left(P(\tilde{e} \to \beta)\varepsilon_{e\alpha} - P(\tilde{e} \to \alpha)\varepsilon_{e\beta} \right) \mathcal{A} \tag{19}$$
$$-4 M_0 M' \sum_{ijk} \mathrm{Im}(U^*_{\alpha i} U_{\beta i} U^*_{ek} U_{\alpha k} U_{ej} U^*_{\beta j}) \mathrm{Im}(\phi_{ik}\phi_{ij}) \Big) \ .$$

The asymmetry is

$$S_{\alpha\bar{\beta}} = \frac{\Delta R_{\alpha\bar{\beta}}}{\sigma(\tilde{e}\tilde{\bar{e}}) + \sigma(\tilde{\mu}\tilde{\bar{\mu}}) + \sigma(\tilde{\tau}\tilde{\bar{\tau}})} \ , \tag{20}$$

where the summed differential slepton cross-section follows trivially from Eq. (15),

$$d\sigma(\tilde{e}\tilde{\bar{e}}) + d\sigma(\tilde{\mu}\tilde{\bar{\mu}}) + d\sigma(\tilde{\tau}\tilde{\bar{\tau}}) = K \left(3|M_0|^2 + |M'|^2 + 2 M_0 M' \right) d\cos\theta \ .$$

As expected, setting $\alpha = \mu$ and summing β reproduces the uncorrelated asymmetry in Eq.(11). Fig. 3 shows a comparison of CP conjugate events $\mu^- e^+$ (solid) and $\mu^+ e^-$ (dashed) due to slepton pair production at the NLC with $\sqrt{s} = 500$ GeV.

Discussion

Our study can be generalized to other processes. In the future Large Hadron Collider (LHC), gluinos will be copiously produced, if the gluino mass is about a few hundred GeV. Through the chain of cascade decays, sleptons may occur in the intermediate state and give dilepton events. With enough CPV slepton flavor oscillation, asymmetries of the type discussed here might be observable. In the case of single gluino production, the single lepton asymmetry in Eq. (6) applies, where $\tilde{\alpha}$ denotes the first slepton produced in the decay chain, and $\tilde{\beta}$ is the second slepton. To avoid washing out of the asymmetry, some reconstruction of the cascade decay would be necessary to at least statistically identify the primary and secondary lepton decay products.

We also note that it is possible the NLC could be operated in a $\gamma\gamma$ mode[13]. In this case, all single lepton asymmetries S_l vanish, but $S_{\alpha\bar{\beta}}$ could be measured, and is given by Eq. (14) or Eq. (20) with M' set to zero. The signal and background rates for slepton pair production are similar to those for intermediate mass charged Higgs production, which are roughly equal after mild cuts[14]. Both the t-channel dominated limit of the e^+e^- collider asymmetry S_α and the dilepton asymmetry measured by a $\gamma\gamma$ collider are proportional to X_{CP}, the respective mass-splitting factors are $\mathrm{Im}(\phi_{12} + \phi_{23} + \phi_{31})$ and $\mathrm{Im}(\phi_{12}^2 + \phi_{23}^2 + \phi_{32}^2)$. Measurement of both asymmetries could help indirectly measure the slepton mass-splitting. Finally, we note that if left-handed polarization of the electron beam is possible, the e^+e^- collider measurement of $S_{\alpha\bar{\beta}}$ would yield the same information as that of the $\gamma\gamma$ collider, since $M'(e^-_L e^+_R \to \tilde{l}_R \, \tilde{\bar{l}}_R)$ vanishes at the tree-level.

Acknowledgment

This work was supported in part by U.S. Department of Energy under Grant number DE-FG02-84ER40173.

Figure Captions

Fig. 1 Dependence of the slepton pair production cross-section on the slepton mass $\bar{m}$, at the Next Linear e^+e^- Collider, with $\sqrt{s} = 500$ GeV. The production of the selectron pair $\tilde{e}_R\tilde{\bar{e}}_R$ (solid curves) depends on the neutralino couplings and masses. Here we assume the lightest neutralino of mass M_χ is purely Bino. We also show $\sigma(e^-e^+ \to \tilde{\mu}_R\tilde{\bar{\mu}}_R)$ (dashed). Note that $\sigma(e^-e^+ \to \tilde{\mu}_R\tilde{\bar{\mu}}_R) = \sigma(e^-e^+ \to \tilde{\tau}_R\tilde{\bar{\tau}}_R)$.

Fig. 2 Uncorrelated asymmetry $-(S_\mu - S_\tau)$ versus $\Delta m/\Gamma$ for $X_{\mathrm{CP}} = X_{\mathrm{CP}}^{\mathrm{Max}}$ (solid) or $X_{\mathrm{CP}}^{\mathrm{Max}} = 0.1 \times X_{\mathrm{CP}}^{\mathrm{Max}}$ (dashed), for $f = 0.75$ and $\Delta m = \Delta m_{12} = \Delta m_{23}$.

Fig. 3 Comparison of CP conjugate events $\mu^- e^+$ (solid) and $\mu^+ e^-$ (dashed) due to slepton pair production at the NLC with $\sqrt{s} = 500$ GeV. We choose the case of maximal mixings, $\theta_1 = \theta_2 = \theta_3 = \frac{\pi}{4}$ and $\Delta m_{12} = \Delta m_{23} = \Gamma$. We illustrate the scenarios of $M_\chi = 50$ or 100 GeV for $\bar{m} = 150$ GeV. Event rates are shown versus the KM phase δ. For $-\pi < \delta < 0$, the event rates can be read by replacing δ by $|\delta|$ and reversing the labels between $\mu^- e^+$ and $\mu^+ e^-$.

Note added: While preparing this manuscript, we learned[15] that the authors of Ref. [4] were extending their work to include analysis of lepton flavor violation in the three-generation case, as well as the CPV effects considered here[16].

References

[1] Particle Data Group, Review of Particles and Fields, Phys. Rev. **D54**, (1996).

[2] F. Gabbiani, E. Gabrielli, A. Masiero, and L. Silvestrini, Nucl.Phys. **B477**, 321 (1996), and references therein.

[3] N. V. Krashnikov, Mod. Phys. Lett **A9**, 2825 (1994);
N. V. Krashnikov, Phys.Lett. **B388**, 783-787 (1996).

[4] N. Arkani-Hamed, H.C. Cheng, J. Feng, and L. Hall, Phys. Rev. Lett. **77**, 1937 (1996).

[5] A study of squark oscillation and CPV effects similar to those discussed here can be found in M. Nowakowski and A. Pilaftsis, Phys. Lett. **B245**, 185 (1990).

[6] V. Barger, K. Whisnant, R.J.N. Phillips, Phys. Rev. Lett. **45**, 2084 (1980).

[7] See Ref. [2] for discussion of strong constraints on left-right slepton mixing.

[8] L.-L. Chau and W.–Y. Keung, Phys. Rev. Lett. **53**, 1802 (1984);
C. Jarskog and R. Stora, Phys. Lett. **208B**, 268 (1988).

[9] E. Commins *et al.*, Phys. Rev. A **50**, 2960 (1994).

[10] J. Ellis, S. Farrara, and D.V. Nanopoulos, Phys. Lett. **114B**, 231 (1982); W. Buchmüller and D. Wyler, Phys. Lett. **121B**, 321 (1983); F. del Aguila, M.B. Gavela, J.A. Grifols, and A. Méndez, Phys. Lett. **126B**, 71 (1983).

[11] R. Becker and C. Vander Velde, in *Proceedings of the European Meeting of the Working Groups on Physics and Experiments at Linear* e^+e^- *Colliders*, ed. P. M. Zerwas, Report No. DESY–93–123C, p. 457.

[12] M. Nojiri, K. Fujii, T. Tsukamoto, Phys. Rev. D **54**, 6756 (1996).

[13] I. Ginzburg, G. Kotkin, V. Serbo, and V. Telnov, Nucl. Instrum. Methods 205, 47 (1983); *ibid*, 210, 5 (1983);
V. Telnov, Nucl. Instrum. Methods **A 294**, 72 (1990).

[14] D. Bowser-Chao, K. Cheung, and S. Thomas, Phys. Lett **B315**, 399 (1993).

[15] A talk given by J. Feng on "*SUSY Flavor and CP Violation at Colliders*" at the Phenomenology Meeting, Madison, Wisconsin, March 17, 1997.

[16] N. Arkani-Hamed, H.C. Cheng, J. Feng, and L. Hall, LBNL-40147, UCB-PTH-97/16, Fermilab-PUB-97/078-T, hep-ph/9704205.

Fig. 1


$\sqrt{s}$ = 500 GeV
M_χ=50 GeV
α=e
M_χ=100 GeV
$\alpha=\mu(\tau)$
σ ($e^-e^+ \rightarrow \tilde{l}_{\alpha_R}\,\bar{\tilde{l}}_{\alpha_R}$) (pb)
1.00
0.50
0.20
0.10
0.05
0.02
0.01
100
125
150
175
200
225
250
$\bar{m}$ (GeV)

Fig. 2



0.4
0.3
0.2
0.1
0.0
$-(S_\mu - S_\tau)$
f=0.75
$X_{CP}=X_{CP}^{Max}=0.096$
$X_{CP}=0.1\times X_{CP}^{Max}=0.0096$
0.1
0.2
0.5
1.0
2.0
5.0
10.0
$\Delta m/\Gamma$

Fig. 3



σ (e$^-$e$^+$ → $\mu^\mp$e$^\pm$ + neutralinos)(pb)
0.150
0.125
0.100
0.075
0.050
0.025
0.000
0
50
100
150
δ (degree)
$\sqrt{s}$ = 500 GeV
$\bar{m}$=150 GeV
$\theta_1=\theta_2=\theta_3=\pi/4$
$\Delta m_{12}=\Delta m_{23}=\Gamma$
M_χ (GeV)
50
100
50
100
μ^+e^-
μ^-e^+